

Table 1 Species composition, occurrence and size range of elasmobranch species examined during the present study

Sr. No.	Species	N	Frequency of occurrence (%)	Size range (cm)	Lm (cm)*	Juveniles (n)	Adults (n)
1	<i>Chiloscyllium griseum</i> (Müller & Henle, 1838)	62	16.00	11 – 581	30.6–55.01	31	31
2	<i>Scoliodon laticaudus</i> (Müller & Henle, 1838)	10	15.00	15–571	32.6–58.41	84	24
3	<i>Himantura walga</i> (Müller & Henle, 1841)	63	10.00	5–372	19.8–35.42	45	18
4	<i>Aetobatus flagellum</i> (Bloch & Schneider, 1801)	10	0.05	20–1052	29.9–53.72	06	04
5	<i>Himantura gerrardi</i> (Gmelin, 1789)	07	0.03	9–282	73.8–132.4	07	00
6	<i>Glaucostegus granulatus</i> (Cuvier, 1829)	06	0.03	9–251	99.3–128.2	06	00
7	<i>Rhinobatus obtusus</i> Müller & Henle, 1841	05	0.04	20–301	37.5–67.31	03	02
8	<i>Himantura uarnak</i> (Gmelin, 1789)	03	0.02	22–352	73.8–132.4	03	00
9	<i>Pastinachus sephen</i> (Forsk. ã, 1775)	01	0.01	352	68.2–1242	01	00
10	<i>Neotrygon kuhlii</i> (Müller & Henle, 1841)	01	0.01	142	29.2–52.42	01	00

Note: 1 total length 2 disc width