

Table 1 Antibacterial activity of tissue extract of *H. pugilinus* against clinically isolated human pathogen (Mean \pm SD)

Organism	Zone of Incubation(mm)						
	Ethyl acetate	Chloroform	Petroleum ether	Methanol	Ethanol	Acetone	Aceto nitrile
<i>A.hydrophila</i>	16.00 \pm 0.54 [*]	4.00 \pm 0.55 [*]	-	6.00 \pm 0.46 [*]	-	-	8.00 \pm 0.54 [*]
<i>E.coil</i>	15.00 \pm 0.63 ^{**}	8.00 \pm 0.48 [*]	-	4.00 \pm 0.87 ^{**}	13.00 \pm 0.72 [*]	10.00 \pm 0.49 ^{**}	13.00 \pm 0.59 [*]
<i>S.typhi</i>	10.00 \pm 0.42 [*]	-	-	-	-	4.00 \pm 0.70 [*]	11.00 \pm 0.52 [*]
<i>S.paratyphi A</i>	21.00 \pm 0.74 [*]	10.00 \pm 0.53 [*]	-	-	17.00 \pm 0.46 [*]	4.00 \pm 0.71 ^{**}	16.00 \pm 0.46 ^{**}
<i>V.cholerae</i>	17.00 \pm 0.55 [*]	-	-	7.00 \pm 0.47 ^{***}	12.00 \pm 0.64 [*]	-	13.00 \pm 0.47 [*]
<i>V.paraheamolyticus</i>	14.00 \pm 0.47 [*]	10.00 \pm 0.54 ^{**}	-	6.00 \pm 0.57 [*]	16.00 \pm 0.45 [*]	4.00 \pm 0.54 [*]	18.00 \pm 0.84 [*]

Note: *Values in the same row with different superscripts are significantly different (P<0.05)