

Table 1 Trophic levels of fish types landed by bait fishermen at the Mida creek

Feeding Guild	Family	Scientific name	Frequency	Trophic Level
1 ^o Carnivores	Gerridae	<i>Gerres oyena</i>	5.4	2.29±0.2
	Lethrinidae	<i>Lethrinus nebulosus</i>	4.5	2.43±0.2
	Ephippidae	<i>Platax orbicularis</i>	2.7	2.46±0.3
	Scaridae	<i>Scarus</i> sp.	0.9	2.50±0.5
	Gobidae	<i>Gobuis keiensis</i>	0.9	2.50±0.5
	Labridae	<i>Coris cuvieri</i>	7.2	2.57±0.2
	Gerridae	<i>G. longilostris</i>	0.9	2.80±0.5
	Lutjanidae	<i>Lutjanus fulviflammus</i>	6.3	2.85±0.2
	Lethrinidae	<i>Lethrinus lentjan</i>	5.4	2.87±0.2
	Lethrinidae	<i>Lethrinus harak</i>	25.2	2.90±0.1
	Haemulidae	<i>Plectorhincus vittatus</i>	1.8	2.93±0.3
	Lutjanidae	<i>Lutjanus argentimaculus</i>	1.8	3.00±0.3
	Gerridae	<i>G. filamentosus</i>	11.7	3.04±0.1
	2 ^o Carnivores	Carangidae	<i>Trachinotes baiboni</i>	0.9
Serranidae		<i>Epinephelus malabaricus</i>	5.4	3.34±0.2
Carangidae		<i>Caranx ignobilis</i>	16.2	3.42±0.1
Drepaneidae		<i>Drepane punctata</i>	0.9	3.50±0.5
Lutjanidae		<i>Terapon jarbua</i>	1.8	3.50±0.3