

Table 1 Length and weight and seasonally distribution of *K. pelamis* samples

Years	Seasons	N	Length (cm) (X \pm SD)	Weight (g) (X \pm SD)
2006-2007	autumn	16	58 \pm 2	1800 \pm 239
	winter	14	59 \pm 2	1800 \pm 255
	spring	16	60 \pm 2	2100 \pm 239
	summer	20	59 \pm 3	2200 \pm 300
2007-2008	autumn	14	56 \pm 3	1750 \pm 295
	winter	16	57 \pm 3	1900 \pm 290
	spring	12	60 \pm 3	2400 \pm 337
	summer	18	62 \pm 3	2750 \pm 342
2008-2009	autumn	16	55 \pm 2	1800 \pm 198
	winter	18	58 \pm 2	2200 \pm 200
	spring	14	60 \pm 2	2350 \pm 291
	summer	23	61 \pm 2	2800 \pm 295
2009-2010	autumn	12	57 \pm 1	1800 \pm 175
	winter	18	59 \pm 2	2200 \pm 252
	spring	16	59 \pm 2	2350 \pm 287
	summer	18	62 \pm 2	2700 \pm 281
2010-2011	autumn	14	58 \pm 2	1950 \pm 289
	winter	18	61 \pm 3	2300 \pm 357
	spring	12	61 \pm 3	2700 \pm 483
	summer	20	65 \pm 2	3200 \pm 309

Note: X=mean, SD= standard deviation, N= Number of samples