

Table 1 Effect of Ecoclean on pH, salinity, temperature and dissolved oxygen in infected shrimp ponds at Varadarajapuram

S.N	DOC	pH	Salinity (PSU)	Temperature (°C)	DO	Remarks
1	56	8.590±0.2	6.684±1.1	29.572±0.5	4.628±0.5	Before WG and WF infection 80 kg/day feed was given
2	57	8.521±0.2	6.732±1.0	29.372±0.5	4.273±0.3	Before WG and WF infection 80 kg/day feed was given
3	58	8.099±0.2	7.817±1.1	29.114±0.7	3.629±0.8	After WG and WF infected 60 kg/day feed was given
4	59	7.796±0.2	8.885±1.0	29.909±1.1	3.641±0.7	5 g of Ecoclean applied per kg feed 2 meals/day 60 kg feed was given
5	60	7.515±0.2	7.456±1.0	30.448±1.2	3.933±0.5	2 meals/day 60 kg/day feed was given
6	61	7.655±0.2	7.504±1.1	30.191±1.1	4.021±0.5	2 meals/day 60 kg/day feed was given
7	62	7.910±0.2	8.477±1.0	30.433±1.1	4.681±0.3	2 meals/day 66 kg/day feed was given
8	63	8.396±0.2	7.880±1.0	29.845±0.9	5.153±0.6	2 meals/day 70 kg/day feed was given
9	64	8.507±0.2	8.761±1.0	29.159±1.1	5.227±0.4	1 meal/day 74 kg/day feed was given
10	65	8.638±0.2	8.682±1.1	29.247±1.0	5.288±0.5	After recovered of WG and WF 80kg/day feed was given

Note: Data were expressed as mean values of duplicate independent experiments with standard errors