

Table 2 Production and specific growth rates of different groups of zooplankton

Traitements	Rotifer (<i>Brachionus sp</i> and <i>Asplanchna sp.</i>)			Cladoceran (<i>Moina sp.</i> and <i>Daphnia sp.</i>)			Copepods (<i>Thermocyclops sp.</i>)		
	P (ind/j)	μ (j ⁻¹)	R ²	P (ind/j)	μ (j ⁻¹)	R ²	P (ind/j)	μ (j ⁻¹)	R ²
T ₀	2±0.48	0.10f	0.78	7±1.54	0.00ef	-0.10	7±11.54	0.00g	0.00
T ₁	108±0.80	0.34e	0.90	13±1.54	0.00e	0.00	87±30.55	0.34f	0.99
T ₂	509±1.27	1.48a	0.94	840±20	1.53a	0.82	260±20	1.28a	0.91
T ₃	455±0.83	0.94c	0.87	153±30.55	1.09c	0.88	193±30.55	1.19b	0.92
T ₄	451±1.27	1.30b	0.81	340±20	1.47b	0.97	220±20	0.89c	0.94
T ₅	253±0.83	0.81d	0.84	180±20	0.54d	0.97	227±29.03	0.89ce	0.98

Note: μ = specific growth rate, R² = coefficient of correlation, P = daily production. Means followed by the same letter are not significantly different (P>0.05) according to the LSD test of Fisher