

Table 2 Distribution of octolasmids attached on the gills of *Portunus sanguinolentus*

Gill No.	Left Branchial Chamber (LBC)								Total in LBC	Right Branchial Chamber (RBC)								Total in RBC	G. Total LBC+R BC	%
	Epibranchial				Hypobranchial					Epibranchial				Hypobranchial						
	P	M	D	T	P	M	D	T		P	M	D	T	P	M	D	T			
1	0	0	0	0	0	4	0	4	4	0	0	0	0	0	2	0	2	2	6	0.12
2	0	0	0	0	15	18	3	36	36	0	0	0	0	23	38	5	66	66	102	2.09
3	2	5	0	7	82	122	33	237	244	2	6	3	11	101	122	34	257	268	512	10.52
4	4	14	2	20	152	230	35	417	437	8	17	4	29	202	275	50	527	556	993	20.40
5	7	17	3	27	179	205	60	444	471	8	22	3	33	171	239	41	451	484	955	19.62
6	5	14	3	22	185	221	69	475	497	6	23	10	39	195	195	69	459	498	995	20.44
7	0	17	2	19	164	187	51	402	421	2	15	3	20	154	151	37	342	362	783	16.09
8	2	10	2	14	130	147	31	308	322	3	6	2	11	77	92	19	188	199	521	10.70
Total	20	77	12	10	907	1134	282	2323	2432	29	89	25	143	923	1114	255	2292	2435	4867	100

Note: P = proximal, M = medial, D = distal