

Table 2 The list of fouling macroalgae of *R. venosa* and their occurrence (0 – taxa is not detected, 1 – rare, 2 – seldom, 3 – common, 4 – very common) within the biotopes of sandy (I) and rocky (II) substrates

Taxa	Occurrence		Depth (m)
	I	II	
Green Algae, Chlorophyta			
<i>Bolbocoleon piliferum</i> Pringsheim	1	0	3–10
<i>Bryopsis plumosa</i> (Hudson) C. Agardh	1	0	3–6
<i>Bryopsis</i> sp.	1	0	3–6
<i>Chaetomorpha aerea</i> (Dillwyn) Kützing*	3	2	1.5–15
<i>Cladophora albida</i> (Nees) Kützing*	2	0	4–6
<i>Cladophora laetevirens</i> (Dillwyn) Kützing	2	0	4–6
<i>Cladophora liniformis</i> Kützing	2	0	2–5
<i>Cladophora sericea</i> (Hudson) Kützing	1	1	1–3
<i>Cladophoropsis membranacea</i> (Hofman Bang ex C. Agardh) Børgesen	3	3	2–15
<i>Codium vermilara</i> (Olivi) Delle Chiaje	1	0	5.0
<i>Phaeophila dendroides</i> (P. Crouan & H. Crouan) Batters	1	0	3–6
<i>Pringsheimiella scutata</i> (Reinke) Marchewianka	1	0	3–6
<i>Rhizoclonium riparium</i> (Roth) Harvey	1	1	2–6
<i>Ulothrix implexa</i> (Kützing) Kützing	1	0	3–6
<i>Ulva clathrata</i> (Roth) C. Agardh	2	1	2–5
<i>Ulva flexuosa</i> Wulfen*	2	1	1–3
<i>Ulva intestinalis</i> Linnaeus	3	2	0.5–10
<i>Ulva linza</i> Linnaeus	2	1	1–5
<i>Ulva prolifera</i> O.F. Müller	1	0	4–5
<i>Ulva rigida</i> C. Agardh*	4	2	1–20
Brown Algae, Ochrophyta (Phaeophyceae)			
<i>Cladostephus spongiosum</i> (Hudson) C. Agardh*	4	2	2–15
<i>Cystoseira barbata</i> (Stackhouse) C. Agardh*	1	0	2.5–4
<i>Cystoseira crinita</i> Duby	1	0	3.5–5
<i>Cystoseira</i> sp.	1	0	4–6
<i>Dictyota dichotoma</i> (Hudson) J.V. Lamouroux	1	0	2.5–6
<i>Dictyota fasciola</i> (Roth) J.V. Lamouroux	1	0	6–7
<i>Dictyota spiralis</i> Montagne	1	0	4–6
<i>Dilophus</i> sp.	1	0	4–6
<i>Ectocarpus siliculosus</i> (Dillwyn) Lyngbye	1	0	6–8
<i>Ectocarpus</i> sp.	1	0	3–6
<i>Myrionema balticum</i> (Reinke) Foslie	1	0	3–6
<i>Nereia filiformis</i> (J. Agardh) Zanardini	1	0	4–6
<i>Padina pavonica</i> (Linnaeus) Thivy	1	1	2.5–3
<i>Ralfsia verrucosa</i> (Areschoug) Areschoug	3	2	2–6
<i>Sphacelaria cirrosa</i> (Roth) C. Agardh*	1	0	4–6
<i>Zanardinia typus</i> (Nardo) P. C. Silva*	2	1	2–8
Red Algae, Rhodophyta			
<i>Acrochaetium</i> sp.	1	1	4–5
<i>Antithamnion cruciatum</i> (C. Agardh) Nägeli	1	0	4–5
<i>Apoglossum ruscifolium</i> (Turner) J. Agardh	1	0	4–5
<i>Callithamnion corymbosum</i> (Smith) Lyngbye	1	1	2–15
<i>Ceramium rubrum</i> C. Agardh	1	0	4–6
<i>Chondria capillaris</i> (Hudson) M.J. Wynne	1	0	4–5
<i>Chondria dasyphylla</i> (Woodward) C. Agardh	1	0	2–14
<i>Coccolytus truncatus</i> (Pallas) M.J. Wynne & J.N.Heine	1	0	19.5
<i>Dasya baillouviana</i> (S.G. Gmelin) Montagne	1	0	5–6

Continued Table 2

Taxa	Occurrence		Depth (m)
	I	II	
<i>Dermocorynus dichotomus</i> (J. Agardh) Gargiulo, M. Morabito & Manghisi	1	0	4–5
<i>Ellisolandia elongata</i> (J. Ellis & Solander) K.R. Hind & G.W. Saunders	1	0	6–8
<i>Gelidium crinale</i> (Hare ex Turner) Gaillon	2	0	4–8
<i>Gelidium spinosum</i> (S.G. Gmelin) P.C. Silva	1	1	2–10
<i>Gracilaria dura</i> (C. Agardh) J. Agardh*	1	0	3–10
<i>Hydrolithon farinosum</i> (J.V. Lamouroux) Penrose & Y.M. Chamberlain	1	1	2–6
<i>Jania rubens</i> (Linnaeus) J.V. Lamouroux	1	1	1.5–8
<i>Laurencia coronopus</i> J. Agardh	1	1	2–6
<i>Laurencia obtusa</i> (Hudson) J.V. Lamouroux	1	0	4.0
<i>Lomentaria clavellosa</i> (Lightfoot ex Turner) Gaillon	1	0	4–5
<i>Lomentaria compressa</i> (Kützing) Kylin	1	0	4.5
<i>Peyssonnelia dubyi</i> P. Crouan & H. Crouan*	4	4	0.5–15
<i>Phyllophora crispa</i> (Hudson) P.S. Dixon*	1	0	5–23
<i>Phymatolithon lenormandii</i> (Areschoug) Adey*	1	2	0.5–15
<i>Polysiphonia breviarticulata</i> (C. Agardh) Zanardini	1	0	4–5
<i>Polysiphonia elongata</i> (Hudson) Sprengel*	1	0	2–8
<i>Polysiphonia opaca</i> (C. Agardh) Moris & De Notaris	1	0	4–5
<i>Polysiphonia violacea</i> (Roth) Sprengel	1	0	3–6
<i>Stylonema alsidii</i> (Zanardini) K.M. Drew	1	0	3–6
<i>Vertebrata subulifera</i> (C. Agardh) Kuntze*	1	0	4–10

Note: *species which have been listed in [15]