

Table 2 Mean and standard deviation of the biometrics of the fish samples

Parameters	All samples (n=12) \pm STD	Group A (n=4) \pm STD	Group B (n=4) \pm STD	Group C (n=4) \pm STD
Total length (cm)	28.22 \pm 3.20	24.90 \pm 0.82	27.75 \pm 1.19	32.00 \pm 1.15
standard length (cm)	23.46 \pm 2.85	20.50 \pm 0.58	23.00 \pm 1.08	26.88 \pm 0.85
body weight (g)	451.75 \pm 120.70	315.75 \pm 13.40 ^c	443.50 \pm 19.09 ^b	596.00 \pm 19.54 ^a
fillet weight (g)	148.66 \pm 34.18	113.49 \pm 3.75 ^c	140.53 \pm 4.69 ^b	191.95 \pm 3.65 ^a
remnant weight (g)	188.85 \pm 56.80	125.91 \pm 4.51 ^c	183.36 \pm 8.76 ^b	257.28 \pm 12.94 ^a
waste weight (g)	114.66 \pm 31.05	77.60 \pm 5.86 ^c	119.61 \pm 14.77 ^b	146.78 \pm 3.49 ^a
fillet weight (%)	33.29 \pm 1.90	35.96 \pm 0.71	31.70 \pm 0.40	32.22 \pm 0.43
remnant weight (%)	41.47 \pm 1.33	39.89 \pm 0.27	41.38 \pm 2.22	43.15 \pm 0.79
waste weight (%)	25.24 \pm 1.21	24.16 \pm 0.93	26.92 \pm 2.45	24.64 \pm 0.36

Note: Means with the same letter are not significantly different ($p > 0.05$)