

Table 3 Mean \pm range of Initial Biomass (IB), Final Biomass (FB), Specific Growth Rate (SGR), Food Conversion Rate (FCR), Survival Rate (SR) and Daily Weight Gain (DWG) of *Hoplobatrachus occipitalis* tadpoles reared at different stocking density

Param ètres	D ₁ (5 tadpoles/L)	D ₂ (10 tadpoles/L)	D ₃ (15 tadpoles/L)	D ₄ (20 tadpoles/L)
IB	7.483 \pm 0.386 ^d	16.450 \pm 0.026 ^c	25.356 \pm 0.621 ^b	39.570 \pm 0.320 ^a
FB	23.750 \pm 0.601 ^a	15.93 \pm 0.113 ^b	15.970 \pm 0.762 ^b	9.386 \pm 0.240 ^c
SGR (%/day)	4.220 \pm 0.255 ^a	1.960 \pm 0.326 ^b	2.251 \pm 0.431 ^b	3.683 \pm 0.561 ^a
FCR	1.436 \pm 0.023 ^c	1.810 \pm 0.075 ^b	2.185 \pm 0.218 ^a	2.176 \pm 0.058 ^a
Sr (%)	86.333 \pm 3.333 ^a	66.777 \pm 0.962 ^b	38.333 \pm 1.111 ^c	30.388 \pm 1.734 ^d
dwg (g/day)	0.735 \pm 0.026 ^a	0.254 \pm 0.018 ^b	0.270 \pm 0.071 ^b	0.022 \pm 0.021 ^c

Note: ^{abc} mean values on the same line and not affected by the same letter are significantly different ($p < 0.05$)