

Table 3 The most common algal consort complexes on the sandy (I) and rocky (II) *R. venosa* with indication of the samples quantity / % for the ecomorph

| Algal complex | Samples quantity / % | |
|---|----------------------|-----------|
| | I | II |
| <i>Chaetomorpha aerea</i> – <i>Cladophora laetevirens</i> – <i>Ulva rigida</i> | 34 / 4.1 | 1 / 0.7 |
| <i>Ch. aerea</i> – <i>Peyssonnelia dubyi</i> – <i>U. rigida</i> | 135 / 16.3 | 3 / 2.0 |
| <i>Ch. aerea</i> – <i>Cladophoropsis membranacea</i> – <i>U. rigida</i> – <i>P. dubyi</i> | 26 / 3.1 | - / - |
| <i>Ch. aerea</i> – <i>U. rigida</i> – <i>Cladostephus spongiosum</i> | 39 / 4.7 | 2 / 1.4 |
| <i>Ch. aerea</i> – <i>P. dubyi</i> | 27 / 3.3 | 8 / 5.5 |
| <i>C. laetevirens</i> – <i>Cladophora liniformis</i> | 19 / 2.3 | 2 / 1.4 |
| <i>Cladophora</i> spp. – <i>U. rigida</i> | 24 / 2.9 | 3 / 2.0 |
| <i>C. membranacea</i> – <i>Gelidium crinale</i> – <i>P. dubyi</i> | 28 / 3.4 | 5 / 3.4 |
| <i>C. membranacea</i> – <i>Ralfsia verrucosa</i> – <i>Zanardinia typus</i> | 21 / 2.5 | 4 / 2.8 |
| <i>C. membranacea</i> – <i>R. verrucosa</i> – <i>P. dubyi</i> | 49 / 5.9 | 5 / 3.4 |
| <i>P. dubyi</i> – <i>C. membranacea</i> | 107 / 12.9 | 22 / 15.2 |
| <i>P. dubyi</i> – <i>R. verrucosa</i> – <i>C. spongiosum</i> | 126 / 15.2 | 24 / 16.5 |
| <i>P. dubyi</i> – <i>Phymatolithon lenormandii</i> | 23 / 2.8 | 28 / 19.3 |
| <i>P. dubyi</i> – <i>U. rigida</i> – <i>U. intestinalis</i> | 43 / 5.2 | 8 / 5.5 |
| <i>P. dubyi</i> – <i>Z. typus</i> | 12 / 1.4 | 17 / 11.7 |
| <i>Z. typus</i> – <i>U. rigida</i> – <i>G. crinale</i> | 27 / 3.3 | 3 / 2.0 |