

Table 3 Mean number of fish identified, mean number of species, and mean Shannon-Weaver-weaver Diversity Index per 5 seconds of video from 3 sampled seasons

Season	Zone	Total	sdv	N° sp	sdv	H'	sdv
Autumn	A	23.0	13.7	0.9	0.9	0.14	0.1
	B	0.5	0.6	0.7	0.1	0.06	0.0
	C	8.3	0.0	0.5	0.0	0.00	0.0
	D	0.0	0.0	0.8	0.0	0.22	0.0
	E	0.0	0.0	0.8	0.0	0.08	0.0
Spring	A	30.9	26.1	0.8	0.7	0.21	0.2
	B	13.6	17.4	0.5	0.4	0.12	0.1
	C	9.0	0.0	5.0	0.0	0.39	0.0
	D	5.0	1.7	0.6	0.3	0.02	0.0
	E	27.0	0.0	0.2	0.0	0.01	0.0
Summer	A	36.1	28.4	2.6	2.2	0.81	0.9
	B	11.9	5.9	1.5	2.3	0.02	0.0
	D	8.2	1.6	0.0	0.0	0.00	0.0
	E	1.8	3.6	0.6	0.4	0.15	0.1

Note: A: Central core HCAR; B: Inside central core HCAR; C: Individual HCAR; D: Inside individual HCAR; E: Sandy bottom with *Caulerpa racemosa*